

Comparing Backstories of a Socially Assistive Robot Exercise Buddy for Adolescent Youth

Katelyn Swift-Spong, Cheng K. Fred Wen, Donna Spruijt-Metz, and Maja J Matarić

Abstract—We describe a study that compares two types of backstories, fictional and realistic, of an autonomous Socially Assistive Robot (SAR) exercise buddy in a four-session exercise intervention with overweight adolescents aged 11-14. We find a positive participant response to the robot as an exercise buddy and a trend toward increased intrinsic motivation for physical activity post-intervention. We do not find differences in ratings of the two backstories or between pre- and post-intervention measures of physical activity enjoyment and activity level.

I. INTRODUCTION

In 2007-08, an estimated 34.7% of children aged 6-19 were overweight or obese [1]. Despite the well-established body of literature that demonstrates the protective effects of physical activity (PA) against obesity and diabetes risks among school-age adolescents [2], adolescents engage in an insufficient amount of PA [3] that is less than 6-11 year old children [4]. Adolescent PA engagement exists at various levels (i.e., environmental, interpersonal, and individual) [5], [6], [7], [8], and the importance of social and peer support in prompting PA in adolescents is well established [6], [7]. While positive communication with friends about PA and joint participation are associated with higher levels of physical activity [6], adolescents, especially those who are overweight and obese, are at risk for social interactions, including teasing and stigmatization, that diminish motivation for PA [9].

Physical activity promotion efforts that harness the supportive aspects of peer and social interactions could be effective for this age group. We examined the efficacy of using a Socially Assistive Robot (SAR) as an exercise buddy that provides peer-level support, interaction, and encouragement to adolescents. If such SAR technology could be made affordable and accessible it could aid in addressing peer-level support and encouragement issues while avoiding stigmatization and embarrassment.

We conducted a four-session PA intervention with 11-14 year old overweight adolescents, using an autonomous SAR system, and comparing a fictional to a realistic robot backstory. Within this context, we define the role of the robot as a *buddy* that performs the task co-actively with the user when possible and has a similar ability level at the task as the user. We find that participants responded positively to the robot exercise buddy. In addition, we find a trend towards increased intrinsic motivation for PA post-intervention.

K. Swift-Spong (corresponding author), C. K. F. Wen, D. Spruijt-Metz, and M. J. Matarić are with The University of Southern California, Los Angeles, CA 90089, USA (email: swiftspo, chengkuw, dmetz, matric@usc.edu).

II. RELATED WORK

SAR systems have been used in the role of exercise coaches in a various domains (e.g., elderly [10], [11], stroke rehabilitation [12], and general exercise [13]). However, SAR in *the role of a peer-level buddy* has not received much research attention. Nass et al. found that people treat computers as teammates when there is task interdependence with the computer [14]. Analogously, adolescents may treat robots as peers when they are placed in a buddy role.

Backstory is the means by which the robot is established in the role of peer-level buddy. Backstory research is new in HRI and SAR, but it has been successfully used in several studies. The Roboceptionist characters all had a strong backstory revealed during interactions [15]; a backstory about dragon race training was used in a study focused on children and nutrition [16]. Both studies used fictional backstories, but no comparison of fictional vs. realistic backstories has been done in SAR. Such comparison has been explored with virtual humans. Kang and Gratch found that people with high levels of social anxiety disclosed more to a virtual human counselor with a human backstory than one with a computer backstory [17]. A long-term study by Bickmore et al. found that people engaged in more discussions with a virtual character with a fictional first person backstory than one with a backstory presented in the 3rd person [18]. Because of the physical differences between the robot and adolescents, in this work we hypothesized that participants would react more positively to a SAR exercise buddy with a fictional backstory than one with a realistic backstory.

III. FOCUS GROUPS

We conducted a series of two-session focus groups with a total of 12 participants aged 11-14 years old in three separate groups. The focus group sessions were performed to inform the design of the interaction and centered around aspects such as the idea of using the Nao robot as an exercise buddy, possible backstories for the robot, exercises, motivational messages, and using music during the exercise sessions. The Nao robot was selected for this study because it is the best available robot platform based on its size, safety, cost, and ability to do some exercises. We used ideas and feedback collected from these focus groups to develop and finalize the backstory of the robot, the exercises, and the motivational messages.

Past work has shown that *affective messages*, which focus on the emotional outcome of a behavior, such as how it affects happiness enjoyment, increase physical activity levels in inactive older adolescents more than *instrumental*

messages, which focus on the beneficial or harmful outcomes such as the effects of exercise on health [19]. Consequently, to design the affective messages for the SAR system, we had focus group participants provide feedback on a list of affective messages in four categories – congratulatory, encouraging, feedback, and testimonial. Examples of those affective messages include: “Lets keep up the good work. Exercising will make us feel energetic and happy.” and “I am so happy that we finished that exercise.”

IV. METHODOLOGY

A. Study Design

The study compared two robot backstories using a four-session within-participants study design with 11-14 year-old overweight adolescents. A within-participants design was used because the study did not include enough participants for a between-participants comparison. The backstories were designed to portray the robot as a peer-level exercise buddy. Each participant took part in four exercise sessions, two with each robot backstory. The order of backstory presentation was randomized using block randomization. In both backstory conditions, the robot presented information about its characteristics at analogous interaction points at the beginning, during, and at the end of each exercise session. Four sessions were used so that each backstory would be seen in two exercise sessions in order to minimize novelty. Questionnaire data related to enjoyment of physical activity, perceived stress, and intrinsic motivation to exercise were collected before the first exercise session and after the fourth, and questionnaire data related to perceptions of the robot were collected after the last session with each robot backstory (the second and fourth exercise session).

B. Experiment Setup

The experiment setup is seen in Figure 1. The humanoid robot, a Nao Next Gen V4, was placed on a table so as to be close to or slightly below eye level of the participant. A monitor next to the robot displayed the exercise sequence for the session, including the name and number of repetitions or duration for each exercise. It was also used to show demonstrations of each exercise. A keypad was placed between the robot and the monitor. Using this keypad participants could respond to questions from the robot using keys labeled “Yes” and “No”, inform the robot of being done with a set of exercises using a key labeled “Done”, and stop or pause the interaction with a key labeled “End”. The robot autonomously responded to these inputs based on a pre-determined script. A 3D Vision (Kinect) sensor hanging from the ceiling and an Augmented Reality (AR) tag on top of the robot’s head were used to capture the position of the robot. The robot used this position information to autonomously move back to its starting position after exercises that involved foot movement. An exercise step and treadmill were also in the exercise area. The robot wore either green or orange wristbands, and the color of these was switched between the two backstories to indicate that the backstory was different. The pre-determined scripts for what the robot with each



Fig. 1. Robot exercise buddy experiment setup. Left: Robot performing a split squat with an arm raise to the side. Right: Robot, step, and treadmill setup. Participants exercised in the space between the table and treadmill.

backstory said were the same except for some of the exercise names and characteristics of the backstory, as described in Table I. In addition, the robot performed scripted non-verbal behaviors such as head nods and shakes, pointing, idle behaviors while talking, and other expressive gestures.

A Kinect sensor was placed behind the robot and monitor and used to record the participant’s body pose information. Two USB cameras and one HD camera were used to record audio and video of the interaction from different angles. The experimenters were in the same room as the participant, but behind an opaque black curtain, in case the participants had any questions or problems, and the participants were aware of the experimenters’ presence behind the curtain.

C. Robot Backstories

Based on the ideas and insights from the focus groups, we developed two robot backstories, one fictional and one realistic. Table I illustrates the differences in how the robot used each backstory to describe itself.

D. Exercise Sessions

The exercise sessions were designed around the idea of circuit training, which has been shown to be effective [20], and included strength training exercises, treadmill, and step up exercises. The exercise sessions were designed to be around 30 minutes long. The exact duration of each session was determined by the participant’s pace in performing the exercises. We chose strength training exercises that the robot and participant could do together. Due to motion limitations, the robot could not perform the following exercises: walking, running on the treadmill, and step ups. The strength training exercises included upper body arm movement exercises such as bicep curls and raising the arms to the front, side, and over the head as well as lower body exercises such as squats, split squats, and side steps. Combinations of upper and lower body exercises, and sequences of two exercises were also included in the circuit training design.

Each session contained three step up exercises (right leg, left leg, and both legs in a quick jumping manner), one five minute walk or run on the treadmill, two sequences of exercises, four exercises that involved the lower body, and two upper body only exercises. Each of the four sessions therefore had a slightly different set of exercises but the total number of exercises was the same across sessions, as was the ratio of upper body, lower body, sequence, treadmill, and

TABLE I
DESCRIPTION OF BOTH ROBOT BACKSTORIES.

Characteristic	Fictional Backstory	Realistic Backstory
Name	Mio	Nao
Identity	I am from another planet called Kepler 186f. I came to Earth because my home planet is unhealthy and I want to prevent Earth from becoming unhealthy too.	I am a robot. There are many robots like me around the world, playing soccer, helping kids, and getting involved in research. I came to Los Angeles to be part of research here at USC.
Reason for exercising	I need to exercise more on Earth because Kepler 186f has low gravity and I need to get stronger in order to live here.	I have been programmed with many exercises, and I am here to share some of my knowledge with you.
Exercise ability	After all, I am from another planet. Kepler 186f has lower gravity than Earth, so my balance isn't so great here.	My motors can only move in certain ways and I am not programmed to do complicated balancing.
Source of demo videos	A friend of mine made videos of these exercises so you will know how to do them.	Researchers at USC gave me some demo videos of these exercises so you will know how to do them.
Speech understanding and vision	I don't see or hear so well because my home planet is super dark and loud.	I've also been programmed to speak English but can not understand it, and I don't have any algorithms to help me see what you are doing.
Source of knowledge	Here is the second set of exercises that I learned during my space travel training on Kepler 186f.	Here is the second set of exercises that I was programmed to do by researchers at USC.

step up exercises. The exercise sessions were presented in the same order for all participants.

To explain how each exercise (except the treadmill) should be done, a pre-recorded demonstration video of a human doing and explaining the exercise was played on the monitor before the exercise began. For the treadmill exercise, an explanation, demonstration, and participant practice of using the treadmill was done by the experimenter before the first exercise session. In addition, participants were allowed to choose the speed of the treadmill and were told that they could either walk or run on the treadmill. Participants were also told that they could do the strength training exercises either at the same pace as the robot or at their own pace. Participants were asked to complete a specified number of repetitions of each of the strength training exercises (10 or 15 depending on the exercise), and they were asked to do each of the treadmill and step up exercises for a certain amount of time displayed using a countdown timer on the monitor.

Participants were instructed to push the “Done” button after completing the repetitions for each strength training exercise, to let the robot know they were finished. The press of the “Done” button was the only input the robot had to know if the participants completed the exercises. As such it would have been possible for participants to press the “Done” button and continue with the session without

actually completing the exercise. After the “Done” button was pushed, the robot gave feedback by saying one of the affective messages described in Section III. The affective messages were pre-scripted and were the same for both robot backstories.

E. Measures

A collection of measures was used, including questionnaires and behavioral measures. Questionnaires administered both pre-intervention (immediately before the first exercise session) and post-intervention (immediately after the fourth exercise session) included a modified version of the *Physical Activity Enjoyment Scale (PACES)* [21], which measures enjoyment of physical activity, a modified version of the *Perceived Stress Visual Analog Scale* [22], which measures perceived stress levels, and the *Exercise Self-Regulation Questionnaire (SRQ-E)* [23], which measures intrinsic motivation for physical activity. In addition, the *Big Five Inventory*, which measures personality traits along five dimensions was given immediately before the first interaction session [24], [25], [26] along with a set of demographics questions. Height and weight (using a Tanita Body Composition Analyzer) were also taken immediately before the first exercise session and after the last exercise session, and BMI was calculated.

Four questionnaires were used to assess the participants’ reactions to the two robot backstories. These were given immediately after the last session for each backstory (the second and fourth exercise sessions). The first three of these questionnaires included the social richness subscale of the *Temple Presence Inventory (TPI)* [27], interaction-related questions modified from the TPI [27], and a modified version of the social attraction subscale of the *Interpersonal Attraction Scale* [28]. Modified versions of these three questionnaires were previously successfully used by Jung and Lee in a study with social robots [29]. The fourth questionnaire was a *Robot Expectations Questionnaire* [30]. In addition, several open-ended questions about the interaction with the robot and the robot’s backstory were asked after the second and fourth sessions. Finally, after the fourth exercise session, participants were also asked an open-ended question about which robot (i.e., backstory) was a more enjoyable exercise buddy and why.

Additional data collected during each exercise session included video, audio, skeletal tracking (using a Kinect), and heart rate (using a Mio Alpha 1 wrist-worn heart rate monitor). Accelerometer data were collected during waking hours (using an Actigraph, Inc. GT3X-BT accelerometer) before the first exercise session and after the fourth exercise session. Analysis of the video, audio, skeletal tracking, heart rate, Perceived Stress Visual Analog Scale, Big Five Inventory, and Robot Expectations Questionnaire data is outside the scope of this paper.

F. Procedure

At the beginning of each exercise session, participants were told that the robot was currently sleeping. They were also told the name of the robot and that they would exercise

with a different robot the following week. Participants were instructed to press the “Yes” button to wake up the robot. When they did so, the autonomous robot interaction began. After participants completed each exercise session and questionnaires, they received compensation of \$20 in cash (an amount typical for the type of study and geographic area).

G. Hypotheses

H1: Participants will have more positive reactions to the robot with a fictional backstory than the robot with a realistic backstory.

H2: Intrinsic motivation for physical activity as measured by the SRQ-E will increase from before the first exercise session to after the fourth exercise session.

H3: Physical activity enjoyment as measured by the PACES will increase from before the first exercise session to after the fourth exercise session.

H4: Activity levels as measured using the ActiGraph activity monitor will increase from before the first exercise session to after the fourth exercise session.

H. Study Population

There were a total of 22 participants, of whom 18 completed the study (Two of the 18 also participated in the focus groups). Participants were recruited from events and organizations in the local area, such as churches and health fairs, and from a participant pool from other health-related studies at USC. Of the four who did not complete the study, two completed the first session only, and the other two completed the first two sessions. Only data from the 18 participants who completed all four sessions were used in the analysis. This set of 18 participants was composed of 12 females and 6 males, with an average age of 12.11 ($SD = 1.28$, $Range : 11 - 15$) and an average Body Mass Index of 27.75 ($SD = 5.54$, $Range : 23.00 - 46.22$) at baseline. Participants were 11-14 years old at the time of recruitment. Fourteen of the 18 participants were self-identified as Hispanic, 2 as mixed ethnicity (1 as Mexican and African American and 1 as African American and French), 1 as White, and 1 as other. The average number of days between the first exercise session and the fourth was 13.17 ($SD = 4.18$, $Range : 10 - 22$). Participants were told that if there were any questions they did not want to answer on the questionnaires, those could be skipped. Thus, some of the questionnaire scales had missing data.

V. RESULTS

A. Evaluation of the Robot with Different Backstories

Using a paired t-test we did not find a statistically significant difference between the participants’ ratings of the fictional and realistic robot backstories when comparing ratings of the social richness ($t(15) = 0.11$, $p = 0.91$; $M_F = 4.85$, $SD_F = 0.72$; $M_R = 4.86$, $SD_R = 0.48$), interaction ($t(-0.07) = 17$, $p = 0.95$; $M_F = 4.82$, $SD_F = 0.94$; $M_R = 4.85$, $SD_R = 1.13$) and social attraction ($t(17) = 0.17$, $p = 0.87$; $M_F = 5.40$, $SD_F = 1.25$; $M_R = 5.33$, $SD_R = 1.17$)

subscales as seen in Figure 2. These ratings were on a seven-point scale, with two participants who did not respond to all the questions used in the social richness subscale rating the fictional backstory.




Fig. 2. Evaluations of the robot with different backstories.

In response to open-ended question about which robot backstory was more enjoyable, 17 participants responded; 7 reported that they either liked both equally or that they were the same, 9 participants chose a single backstory, and one mentioned liking both and also mentioned one of the backstories. Several themes emerged in response to the question of what participants liked about exercising with the robot with each backstory. Some participants enjoyed feeling comfortable and not pressured exercising with the robot. For example, one said: “I feel more comfortable, I can correct my mistake without being shy or embarrassed.” Several participants also mentioned liking that the robot did the exercises along with them and encouraged them to continue doing the exercises. When asked what they disliked, participants responded that the robot could not do all the exercises with them, that it was slow, and that it asked the same questions multiple times.

B. Intrinsic Motivation for Physical Activity

We calculated the subscales of the SRQ-E, as seen in Figure 3, measured on a seven-point scale and listed from most internally and autonomously motivated, on the left, to least, on the right. Several participants did not answer all of the questions for each subscale (one for pre-intervention Intrinsic Motivation, two for post-intervention Intrinsic Motivation, three for pre-intervention Identified Regulation, two for pre-intervention Introjected Regulation, one for post-intervention Introjected Regulation, and two for pre-intervention External Regulation). Using paired t-tests, we did not find any significant differences between the participants’ ratings pre- and post-intervention for any of the subscales: Intrinsic Motivation ($t(14) = -1.64$, $p = 0.12$; $M_{PRE} = 5.32$, $SD_{PRE} = 1.69$; $M_{POST} = 5.59$, $SD_{POST} = 1.11$), Identified Regulation ($t(14) = -0.80$, $p = 0.44$; $M_{PRE} = 5.67$, $SD_{PRE} = 1.60$; $M_{POST} = 5.67$, $SD_{POST} = 1.14$), Introjected Regulation ($t(14) = 0.93$, $p = 0.37$; $M_{PRE} = 3.25$, $SD_{PRE} = 1.45$; $M_{POST} = 2.87$, $SD_{POST} = 1.49$), and External Regulation ($t(15) = 1.03$, $p = 0.32$; $M_{PRE} = 2.14$, $SD_{PRE} = 0.95$; $M_{POST} = 1.99$, $SD_{POST} = 1.14$). We also calculated the *Relative Autonomy Index (RAI)* as a weighted sum of the four subscales that gives the weights

of 2, 1, 1, and 2 to the Intrinsic Motivation, Identified Regulation, Introjected Regulation, and External Regulation subscales respectively for a maximum rating of +24 and a minimum of -24 [31]. Using a paired t-test, we found an increase in RAI from pre- to post-intervention that was approaching significance ($t(12) = -1.85, p = 0.09; M_{PRE} = 8.95, SD_{PRE} = 5.00; M_{POST} = 9.64, SD_{POST} = 4.92$) with three participants who did not respond to all the questions used to calculate the pre-intervention measure, one for the post-intervention measure, and one for both pre- and post-intervention measures.

C. Physical Activity Enjoyment

Using a paired t-test, we did not find a statistically significant difference between participants' ratings of physical activity enjoyment before the first exercise session and after the fourth, using the PACES measure ($t(14) = 0.97, p = 0.35; M_{PRE} = 4.01, SD_{PRE} = 0.48; M_{POST} = 4.04, SD_{POST} = 0.35$). Participants had high measures of physical activity enjoyment both before and after the intervention on a five-point scale, with one participant who did not respond to all the questions in the pre-intervention measure and two for the post-intervention measure.

D. Activity Levels

Participants were asked to wear the accelerometer around their waist for 24 hours per day (except during activities that involved water) for 7 days both before the first exercise session and after the fourth. The accelerometer was set to collect body movement data for each 30 second epoch.

The raw accelerometer data were downloaded, validated, and processed using ActiLife 6.11. The non-wear time was defined as more than 60 consecutive minutes of zero activity count, with an allowance of 2 minutes of spike between 0 and 100. A valid wear day was defined as at least 600 minutes of wear time. 11 participants and 5 participants had two or more valid wear days in pre- and post-tests, respectively. Sedentary behavior and moderate-to-vigorous physical activity were calculated using cut points described in Evenson et al. [32].

The differences between participants' objectively measured percentage of time spent in 1) sedentary behavior and 2) MVPA pre- and post-intervention were separately evaluated using paired t-tests. There were no significant differences in percentage of time spent in sedentary behavior ($t(2) = 1.07, p = 0.40$) and MVPA ($t(2) = 2.20, p = 0.16$) between pre- and post-intervention accelerometer wear.

VI. DISCUSSION

We did not find a difference between the ratings of the two robot backstories for the social richness, interaction, or social attraction subscales, therefore failing to support the first hypothesis, *H1*. In addition, 7 of the 17 participants who responded to the open-ended question about which robot backstory they enjoyed more liked them both equally or saw them as the same. These results could indicate that the differences between the backstories were not salient enough to change the nature of the interaction. This lack of salience




Fig. 3. Ratings of intrinsic motivation for physical activity.

could in part be due to the fact that several aspects of the robot did not change between the backstories. Specifically, the non-verbal behaviors, the sound of the voice, and the appearance only changed slightly when the color of the wristbands was changed between backstories. This could indicate that backstory differences need to be accompanied by salient changes in robot behavior and appearance in order to be perceived as meaningful. It is also possible that differences could be found without changing the salience if a between-participants study design were used.

Participants had slightly positive (above neutral) reactions to the robot with the two backstories on the social richness, interaction, and social attraction subscales. They also mentioned feeling comfortable exercising with the robot, liking that the robot did exercises along with them, and liking the encouragement from the robot. These positive reactions could indicate that a fully autonomous SAR exercise buddy might be accepted by adolescents.

The second hypothesis, *H2*, was not fully supported by the SQR-E, but there was a trend approaching significance that RAI increased from pre- to post-intervention. Because this robot exercise buddy intervention, however, was not compared to a control with no robot there is no way to know if this change is due to the interaction with the robot.

We did not find support for *H3* as there were no significant differences between physical activity enjoyment ratings before and after the intervention. This could in part be due to a ceiling effect because the physical activity enjoyment ratings before the intervention were already high.

The fourth hypothesis, *H4*, was not supported by pre-post differences in activity levels. This lack of change could be due to the short duration of the intervention. In addition, the intervention did not include strategies or encouragement for participants to continue exercising outside of the experiment sessions. Future work should focus on longer term interventions, and include encouragement for behavior change outside of the intervention sessions.

VII. CONCLUSIONS

This paper described a fully integrated autonomous SAR system capable of serving as a peer-level exercise buddy. The system was evaluated in a study with overweight adolescents that compared fictional and realistic robot backstories. Although we did not find any differences in participants'

ratings of the robot characters, we did find that participants reacted positively to the robot exercise buddy. This suggests that adolescents may be accepting of a robot in the exercise buddy role with a realistic or fictional backstory. In addition, we found a trend toward increasing intrinsic motivation for physical activity after the intervention. This early work provides promising insights toward the development of SAR peer-level robot buddies and calls for further investigation.

ACKNOWLEDGMENTS

This work was supported by the National Science Foundation (NSF) grant IIS-1117279, NSF Graduate Research Fellowship to K. Swift-Spong, and by the National Center for Advancing Translational Sciences of the National Institutes of Health under Award Number UL1TR000130 (formerly by the National Center for Research Resources, Award Number UL1RR031986). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NSF or National Institutes of Health. We thank Jacqueline Kiwata, Professor Todd Schroeder, and Gillian O'Reilly for their help.

REFERENCES

- [1] C. L. Ogden, M. D. Carroll, L. R. Curtin, M. M. Lamb, and K. M. Flegal, "Prevalence of high body mass index in us children and adolescents, 2007-2008," *Jama*, vol. 303, no. 3, pp. 242-249, 2010.
- [2] I. Janssen and A. G. LeBlanc, "Review systematic review of the health benefits of physical activity and fitness in school-aged children and youth," *Int. J. of Behavioral Nutrition and Physical Activity*, vol. 7, no. 40, pp. 1-16, 2010.
- [3] P. C. Hallal, L. B. Andersen, F. C. Bull, R. Guthold, W. Haskell, U. Ekelund, L. P. A. S. W. Group *et al.*, "Global physical activity levels: surveillance progress, pitfalls, and prospects," *The Lancet*, vol. 380, no. 9838, pp. 247-257, 2012.
- [4] R. P. Troiano, D. Berrigan, K. W. Dodd, L. C. Masse, T. Tilert, M. McDowell *et al.*, "Physical activity in the united states measured by accelerometer," *Medicine and Sci. in Sports and Exercise*, vol. 40, no. 1, p. 181, 2008.
- [5] J. F. Sallis, J. J. Prochaska, W. C. Taylor *et al.*, "A review of correlates of physical activity of children and adolescents," *Medicine and Sci. in Sports and Exercise*, vol. 32, no. 5, pp. 963-975, 2000.
- [6] C. C. Maturo and S. A. Cunningham, "Influence of friends on childrens physical activity: a review," *American J. of Public Health*, vol. 103, no. 7, pp. e23-e38, 2013.
- [7] C. Craggs, K. Corder, E. M. van Sluijs, and S. J. Griffin, "Determinants of change in physical activity in children and adolescents: a systematic review," *American J. of Reventive Medicine*, vol. 40, no. 6, pp. 645-658, 2011.
- [8] D. R. Lubans, C. Foster, and S. J. Biddle, "A review of mediators of behavior in interventions to promote physical activity among children and adolescents," *Preventive Medicine*, vol. 47, no. 5, pp. 463-470, 2008.
- [9] P. Rukavina and W. Li, "School physical activity interventions: do not forget about obesity bias," *Obesity Reviews*, vol. 9, no. 1, pp. 67-75, 2008.
- [10] J. Fasola and M. J. Matarić, "Using socially assistive human-robot interaction to motivate physical exercise for older adults," *Proc. of the IEEE*, vol. 100, no. 8, pp. 2512-2526, 2012.
- [11] P. Gadde, H. Kharrazi, H. Patel, and K. F. MacDorman, "Toward monitoring and increasing exercise adherence in older adults by robotic intervention: a proof of concept study," *J. of Robotics*, vol. 2011, 2011.
- [12] K. Swift-Spong, E. Short, E. Wade, and M. J. Matarić, "Effects of comparative feedback from a socially assistive robot on self-efficacy in post-stroke rehabilitation," in *Proc. of the 14th IEEE Int. Conf. on Rehabilitation Robotics (ICORR)*, 2015, pp. 764-769.
- [13] D. Ramgoolam, E. Russell, and A. B. Williams, "Towards a social and mobile humanoid exercise coach," in *Proc. of the 9th ACM/IEEE Int. Conf. on Human-Robot Interaction (HRI)*, 2014, pp. 274-275.
- [14] C. Nass, B. Fogg, and Y. Moon, "Can computers be teammates?" *Int. J. of Human-Computer Studies*, vol. 45, no. 6, pp. 669-678, 1996.
- [15] R. Simmons, M. Makatchev, R. Kirby, M. K. Lee, I. Fanaswala, B. Browning, J. Forlizzi, and M. Sakr, "Believable robot characters," *AI Magazine*, vol. 32, no. 4, pp. 39-52, 2011.
- [16] E. Short, K. Swift-Spong, J. Greczek, A. Ramachandran, A. Litoiu, E. C. Grigore, D. Feil-Seifer, S. Shuster, J. J. Lee, S. Huang *et al.*, "How to train your dragonbot: Socially assistive robots for teaching children about nutrition through play," in *Proc. of the 23rd IEEE Int. Symp. on Robot and Human Interactive Communication (Ro-Man)*, 2014, pp. 924-929.
- [17] S.-H. Kang and J. Gratch, "People like virtual counselors that highly-disclose about themselves," *The Annu. Review of CyberTherapy and Telemedicine*, vol. 167, pp. 143-148, 2011.
- [18] T. Bickmore, D. Schulman, and L. Yin, "Engagement vs. deceit: Virtual humans with human autobiographies," in *In Proc. of the 9th Int. Conf. on Intelligent Virtual Agents (IVA)*, 2009, pp. 6-19.
- [19] R. Sirriyeh, R. Lawton, and J. Ward, "Physical activity and adolescents: an exploratory randomized controlled trial investigating the influence of affective and instrumental text messages," *British J. of Health Psychology*, vol. 15, no. 4, pp. 825-840, 2010.
- [20] J. N. Davis, L. E. Gyllenhammer, A. A. Vanni, M. Meija, A. Tung, E. T. Schroeder, D. Spruijt-Metz, and M. I. Goran, "Startup circuit training program reduces metabolic risk in latino adolescents," *Medicine and Sci. in Sports and Exercise*, vol. 43, no. 11, p. 2195, 2011.
- [21] R. W. Motl, R. K. Dishman, R. Saunders, M. Dowda, G. Felton, and R. R. Pate, "Measuring enjoyment of physical activity in adolescent girls," *American J. of Preventive Medicine*, vol. 21, no. 2, pp. 110-117, 2001.
- [22] S. Cohen, T. Kamarck, and R. Mermelstein, "A global measure of perceived stress," *J. of Health and Social Behavior*, pp. 385-396, 1983.
- [23] R. M. Ryan and J. P. Connell, "Perceived locus of causality and internalization: examining reasons for acting in two domains," *J. of Personality and Social Psychology*, vol. 57, no. 5, p. 749, 1989.
- [24] O. P. John, E. M. Donahue, and R. L. Kentle, "The Big Five Inventory: Versions 4a and 54," *University of California, Berkeley, CA, Institute of Personality and Social Research*, 1991.
- [25] O. P. John, L. P. Naumann, and C. J. Soto, "Paradigm shift to the integrative Big Five Trait taxonomy," in *Handbook of Personality: Theory and Research*, 2008, pp. 114-158.
- [26] V. Benet-Martínez and O. P. John, "Los Cinco Grandes across cultures and ethnic groups: multitrait multimethod analyses of the Big Five in Spanish and English," *J. of personality and Social Psychology*, vol. 75, no. 3, pp. 729-50, sep 1998.
- [27] M. Lombard, T. B. Ditton, D. Crane, B. Davis, G. Gil-Egui, K. Horvath, J. Rossman, and S. Park, "Measuring presence: A literature-based approach to the development of a standardized paper-and-pencil instrument," in *3rd Int. Workshop on Presence*, vol. 240, Delft, The Netherlands, 2000.
- [28] J. McCroskey and T. McCain, "The Measurement of Interpersonal Attraction," *Speech Monographs*, vol. 41, no. 1961, pp. 261-266, 1974.
- [29] Y. Jung and K. Lee, "Effects of physical embodiment on social presence of social robots," in *Proc. of Presence*, 2004, pp. 80-87.
- [30] M. Lohse, "Bridging the gap between users' expectations and system evaluations," in *Proceedings of the 20th IEEE Int. Symp. on Robot and Human Interactive Communication (Ro-Man)*, Atlanta, Georgia, 2011, pp. 485-490.
- [31] W. S. Grolnick and R. M. Ryan, "Parent styles associated with children's self-regulation and competence in school," *J. of Educational Psychology*, vol. 81, no. 2, p. 143, 1989.
- [32] K. R. Evenson, D. J. Catellier, K. Gill, K. S. Ondrak, and R. G. McMurray, "Calibration of two objective measures of physical activity for children," *J. of Sports Sciences*, vol. 26, no. 14, pp. 1557-1565, 2008.