

Using Phone-Based Activity Monitors to Promote Physical Activity in Older Adults: A Pilot Study

Harshvardhan Vathsangam, Gaurav S. Sukhatme

Abstract—Physical inactivity is a leading risk factor in a number of chronic diseases. Older adults are at the highest risk because they are the least inactive of all age groups. Smartphone-based intervention techniques present the opportunity to support cost-effective behavioral intervention to promote physical activity. We present a pilot study that examines the feasibility of utilizing the "Strive" smartphone-based passive sensing application to motivate older adults to stay active. Eight participants aged between 50 and 80 years used the application over a period of three weeks to track their daily activity habits. Participants carried smartphones to track daily activities and introspected about their lifestyles as part of weekly check-in sessions. The value that 24x7 smartphone-based sensors provided to this population was an easy-to-use and always available companion to promote awareness about their physical activity. The main motivations for participants to stay active were to prevent loss of personal health due ageing and the need to stay young. Participants increased their daily physical activity by 15% over the first week. This increase was mainly driven by the younger participants. Participants were willing to bear the inconvenience of physical activity tracking due a phone if it meant that they would obtain a better report of their daily physical activities. Any application that tracks physical activity using smartphones needs to address the wearability of the phone.

Index Terms—Smartphone, Physical Activity, Passive Sensing

I. INTRODUCTION

Physical inactivity and sedentary lifestyles are positively associated with obesity [1], metabolic syndrome [2], type 2 diabetes [3], and mortality [4]. These issues are particularly important for older adults because older adults are the least inactive [5] and account for the highest medical expenditures of any age group in the United States [6]. It is estimated that nearly one out of 5 adults will be 65 and older by 2030. This rapid growth coupled with unhealthy habits has the potential to negatively impact public health.

Lifestyle interventions are one mechanism of addressing health and well-being in older adult populations. However, the cost of implementing lifestyle interventions remains a major challenge behind their widespread adoption [7]. Integrating technology to support such interventions has shown empirical evidence reducing these costs [8]. In particular, smartphone-based technologies represent a unique way to support lifestyle interventions by leveraging hardware that is increasingly ubiquitous in daily life [9]. Smartphones carry a suite of sensors including accelerometers, gyroscopes and GPS units that enable cost-effective, passive and ubiquitous activity monitoring. This in turn allows the delivery of just-in-time, personalized interventions using objective data collected in free living conditions [10].

Previous research has integrated smartphone technology to target a wide range of health conditions and improve health behaviors [11]. In the context of physical activity, there exists a large body of research using mobile phones and behavioral interventions such as goal-setting, social sharing, rewards/badges etc. to promote physical activity

[12, 13, 14, 15]. These approaches have relied on self-reporting based on recall or on reporting of data from external sensors such as pedometers. These approaches could be further improved by combining sensing and intervention in the same platform thus reducing the need for data entry and allowing for more prompt delivery of interventions. Few approaches have used a single, smartphone-based platform that combines on-phone passive sensing with intervention techniques to measure physical activity [16, 17], particularly in older adults [10]. While majority of previous research has focused on the behavioral aspects of such applications, little research has focused on how a smartphone-based passive-sensing tool would be perceived and used by older adults from a technological perspective. Identifying these attitudes could be a key factor in designing future applications that deliver effective physical activity interventions.

This paper seeks to extend previous research by understanding how a smartphone-based, 24x7, passive sensing application can inform older adults about their physical activity and motivate them to be physically active. We use the "Strive" mobile application to sense physical activity. Through a pilot study with eight older adults, we identify design and usability guidelines for the use of this application in this demographic. We explore how such an application could be effectively used to help users gain self-awareness about their activities and increase activity levels. The resultant platform can be used as a foundational tool on which future intervention studies can be built. We also present a detailed survey of phone-usage habits and attitudes toward smartphones among older adults. Through this study, we aim to present a user-centered design approach in a translational setting and lay the groundwork for a larger study in motivating adults to stay physically active.

II. METHODS

A. Demographic Details

The study was designed to run for three weeks and was approved by the Institutional Review Board of the University of Southern California. Participants were recruited from the members of OASIS, a non-profit organization dedicated to help older adults lead productive lives. 16 participants (all female) volunteered of which eight were selected for the study. Participants were divided into groups corresponding to two community centers (Sites A and B) according to their convenience. The average age was 74 ± 11 years (Min: 61, Max: 88). The average weight was 68 ± 15 kg (Min: 52, Max: 99). The average height was 160 ± 7 cm (Min: 150, Max: 170) and average BMI was 26 ± 6 (Min: 22, Max: 40). All participants were retired. Study assistance for recruiting, user support and training was provided by the Front Porch Center for Technology Innovation and Well being.

B. Movement Measurement

Each participant was provided a Google Nexus 4 smartphone with the application pre-installed for the duration




Figure 1: Example application screen showing daily activity summary and weekly activity trends

of the study. Each smartphone ran the Strive application. The Strive application was a custom-built application that passively sensed triaxial accelerometer information at 50 Hz throughout the day through a background process. For each minute of the day, the application would calculate a “Motion Intensity” feature. The Motion Intensity feature was calculated as the sum of the average integrals of the mean-subtracted sensor output per minute across three accelerometer axes. This feature was previously validated against energy expenditure from walking and against traditional accelerometers [18]. This feature was calibrated with a binary classifier to determine whether the user was performing Moderate to Vigorous Intensity Physical Activity (MVPA) or at rest. The application thus provided a minute by minute report of whether the participant was engaging in MVPA. MVPA was chosen because of its relevance in reducing sedentary time [19]. To improve the robustness of the system, an additional rule was implemented stating that if a person were at rest, there must be at least two consecutive minutes of MVPA for the application to decide that the person is moving. This rule was implemented to encourage sustained bouts of physical activity and to avoid false positives. No GPS data were collected. The application automatically tabulated measured active minutes towards a fixed goal. For the purposes of this study, the goal was set at 30 minutes of activity per day based on previously established guidelines [20]. Participants could open the application at any time to check how many minutes of activity were collected and whether they met their goal.

C. Application Design

The design of the application was based on previous work related to usability guidelines for seniors [21]. It consisted of a single home screen with the total active minutes of each day displayed and no menus to avoid navigation hierarchies. The application itself could be directly accessed through the phone home screen through a large widget. Within the home screen, a large, circular progress bar with accumulated active minutes in the center was chosen to indicate goal progress. To show weekly trends, a two-level bar chart was used indicating whether a person was below or above their daily goal. This chart would reset every week.

The design featured just three colors with high contrast and a flat texture. Blue and green were chosen as base colors to stay in the middle of the visible color spectrum. Blue was

used as a base color, green was used to indicate progress. A third color, yellow was used to indicate achievement of goals beyond progress. The application had two color modes: prior to goal completion, the daily activity circle would be blue with a green progress bar; upon completion of goals, the activity circle would be green (to indicate goal achievement) with a yellow completed ring to indicate that all future activity is above the set goal. This allowed a user to quickly estimate how far they were from their goals. Font-sizes were specifically chosen to be large and easily visible from a distance.


D. Study Details

1) *Pre-study orientation:* Prior to commencement of the study, participants attended an orientation session at each center. During each orientation session, participants were given training on how to use the Nexus 4 smartphone and the Strive application. Each participant was told how the application would passively collect their physical activity information and that in order to accurately log their physical activity, they needed to carry their phone on their person for as long as possible. Participants then participated in a demonstration walk where they walked around the center environs with the phone on their person and observed how the application added active minutes based on how long they walked. Participants were allowed to meet and collaborate with other participants from the same center if they felt it would boost their activity levels. Two days after their initial orientation, study volunteers called up the participants to verify that there were no adverse events.


2) *Weekly check-ins:* At the end of each week, the study coordinator met the participants at each center to manually download the information collected by the application. Participants were not required to meet at the same time. The data from each participant was processed using custom software built in Matlab to display relevant summaries of activity. The summary data included the total active minutes they had accumulated each day since the beginning of the study, the total active minutes per week, their weekly summarized activity information and an activity intensity map indicating when they were most active throughout the week. Each participant was also told their ranking (of total minutes accumulated) in their center and across all participants. The coordinator also collected any feedback about the application.

When displaying summaries of data, the study coordinator guided the participants through their data. This guidance included showing the participants at what times they had been more active and asking them to introspect on what they did during those times. This context allowed participants to understand the overall picture behind their activity and when they gained the most active minutes.

3) *Post-study debrief:* At the end of the third week, five out of eight participants attended a post-study focus group. The aim of the focus group was to determine the attitudes and responses of the participants to the application and to smartphone usage in general. Participants filled out a post-study survey that collected information about application usability, wearability and engaged in a group interview with the study coordinator. At the end of the focus group, each participant received a printed summary of their activity over three weeks, activity patterns over time and their ranking (of



(a) Accumulated active minutes per each week of the study



(b) Per-participant cumulative active minutes as the study progressed

Figure 2: Illustration of total accumulated minutes over the study period. Overall participants showed a 15% increase over the first week. This main contributions were from the Site B where participants were younger.

total active minutes accumulated) in their center and across all participants.

III. RESULTS AND DISCUSSION

A. Pre-Study Surveys

1) *Phone habits*: Participants were either first time or novice (less than three months since purchase) smartphone users. Three out of the eight participants were current smartphone users (all iPhone users). Smartphone participants mainly used their smartphones for phone calls, entertainment, shopping, banking, paying bills, accessing government services, and navigation (e.g., looking up a location or route). The remaining five participants used feature phones.

The most common location where participants kept their phones was their shoulder purses or hand purses. The shoulder purse location was the dominant location when the participants were at home, driving and when walking. Other locations included on a bookshelf, tables at home or in bra/fanny pack. At nights, all participants placed their phone on a table or nightstand. The main reasons that drove their decision behind phone placement was convenience and accessibility. This was a consistent reason provided in both home and outdoor settings. Other reasons included where they were least likely to forget and being close to a power supply. The least important reason was the safety of the phone. The fact that the shoulder purse is such a commonly used location implies that any algorithm that tracks user movements for this demographic has to be validated for accuracy in this location.

2) *Motivation for participation in the study*: The main motivational factors for the participants in this study were the need to fight aging, feel good about themselves and maintain overall health. Specific reasons included the need to maintain weight or manage conditions such as osteoporosis and arthritis. One participant saw how not being physically active adversely affected people later in their lives. This made her want to stay active. Other motivations included the need to be active outside of their social commitments and for self-awareness (to understand how busy or not busy they were all day). The least popular reasons were to have fun and

to have a challenge of staying active. Participants preferred low impact activities like walking or swimming for exercise. This suggests that intervention techniques for this age group could focus on creating low-impact exercise suggestions for higher compliance. The messages in these interventions that are most likely to motivate participants to stay active is how exercising helps them stay young and helps them feel good about themselves and less about competitive aspects of activity tracking.

B. Objective Activity Statistics

1) *Summary of Accumulated Activity*: Our system allowed the capture of objective activity information in free-living settings. As part of this study, we collected a total of 241,920 minutes of activity information. As a result, we were able to obtain detailed information about each participant's activity lifestyles. Figure 2 describes the total activity across participants. Participants accumulated a total of 323 hours, 19 minutes over three weeks. Participants from site A logged a total of 193 hours, 35 minutes while those from site B logged a total of 131 hours, 44 minutes. In both sites, a single participant accounted for more than a third of all accumulated minutes (78 hours, 49 minutes for Site A and 107 hours, 51 minutes for Site B). Four out of eight participants accumulated approximately 1100 minutes or 52 minutes of activity in a day. Overall, participants showed a 15% increase over baseline activity in the first week. However this increase came exclusively from Site B participants. One reason for higher minutes could be that Site B participants were on average younger than those from Site A and hence had higher activity capability. Another reason could be that Site B participants were more adept at using the app to gain awareness and modify their lifestyles.

2) *Weekly Activity Patterns*: Figure 3 illustrates the average active minutes across all participants for each day of the study. To examine if there were any patterns in behavior, the days corresponding to the lowest active minutes per day were marked. These days corresponded to the weekends (Saturday/Sunday). This indicates a periodicity in activity patterns. Participants felt that weekends were when they could relax after maintaining high activity during the rest




Figure 3: Illustration of total accumulated minutes per day across all participants. The lowest values in this series corresponded to weekend days (Saturday or Sunday) indicating periodic patterns in activity behavior.




Figure 4: Illustration of typical weekly activity heat maps for a moderately active person in the study. Such maps allowed participants to introspect on their daily activities and identify patterns in their lifestyles.

of the week. Utilizing such patterns would allow evidence-based modification of physical activity interventions. For example, if it were important that participants maintain a consistent activity pattern throughout the week, interventions could be tailored to engage participants to increase their motivation and activity levels during their weekends.

3) *Activity Intensity Maps*: Each participant had the opportunity to view intensity maps of their weekly activity between check-ins. The intensity map consisted of an hourly summary of total active minutes for each day of the week. The purpose behind viewing these intensity maps was for them to reflect on their past behavior and to put their weekly activity in context by identifying patterns. Figure 4 illustrates an activity intensity map for a participant with moderate levels of physical activity. Darker colors indicate higher activity. This intensity map shows that the participant had regular bouts of activity between 10 am and noon every




Figure 5: A strong correlation was observed between accumulated active minutes and age. Younger participants accumulated more minutes. However, the effect tapered at higher age levels.

day of the week (as shown in black). Depending on the intervention needs, the participant could either build on this daily habit or start new habits such as being active in the evening as well to complement their morning activities. Such maps would also allow identification of abnormalities in behavior. For example, one participant fell ill during the course of the study and as a result, their intensity map indicated low levels of activity for five days at a stretch. Identifying such abnormalities would allow timely intervention to maintain health.

4) *Relationship between activity levels and age*: Figure 5 illustrates the relationship between accumulated active minutes in the three week period as a function of age. A quadratic line of best fit was applied. Accumulated minutes decreased with age. All participants carried their smartphones as much as possible, hence this phenomenon could not be explained by lack of wearing a smartphone alone. One reason for this could be that older adults on average have lower physical ability to stay active. Another reason could be that while all adults carried their phones equally, the younger adults were more motivated by the idea of using smartphones to keep track of their activities. We aim to explore this relationship in future work to see if it exists in larger populations. If so, this finding has implications for setting age-appropriate goals of physical activity.

C. Post-Study Surveys

1) *Application Utility*: The most important utility that participants found from the study was using the application data to gain awareness about their lifestyles. Participants checked their activity data whenever they performed large bouts of activity or at specific times of the day such as the afternoon or before going to bed. These habits have implications for fine-tuning the design of interactions between the application and the user. Participants felt that the application was something they could always check, since it was with them all the time and monitoring activity in the background. While the most important feature was using the application data to gain awareness, the least important feature was looking at the application data alone. The second

most important feature was talking about the application data with the study coordinator. This suggests that it is not simply the data that is important but the application plus some contextual insight about lifestyles that matters to this demographic and the ability to interpret the data is important. This has implication on how these data are presented so as to provide contextual insights. Introspection about the context behind activities allowed participants to better understand the reasons behind their variations in activities and modify lifestyles accordingly. For example, one participant was active above average for her grandchild's birthday understood why she was not as active the next day (because of pain in her knee from moving too much).

Goal setting was considered a very important feature of the application. Even though the application assumed a standard goal of 30 minutes per day, all participants set customized goals mentally and tried to meet them. Participants were disappointed if they did not meet their goals. Post-study surveys suggested that allowing participants to set customized goals would help them personalize their experiences and challenge themselves further. This counters what participants felt was the least popular reason to stay active: the need to challenge themselves.

2) *Usability*: Strive was considered simpler to use than traditional pedometers by all participants because no calibration was necessary. The usage of number of active minutes as a unit was considered more intuitive than step counts by all participants. Participants felt that with a traditional pedometer, one has to align steps and that calibration is difficult. Participants felt that the use of active minutes simplified their cognitive load since almost any kind of movement they did could count as activity. This in turn expanded the range of activities that they could consider as exercise. The use of a central number that was easy to access greatly simplified their experience. The support for a no-calibration approach stresses the importance of balancing the need for sensing accuracy with convenience and usability.

3) *Addressing Phone Wearability*: Most of the suggested improvements centered on improving the wearability of the phone. One participant wanted a smaller wearable monitor that would interface with the application that could be clipped and also worked for swimming. She wanted clips or lanyards to extend the range of places where she could wear the phone. Two participants expressed disappointment that the phone did not always pick up their active minutes. One participant noted that when using her stationary bike, the application did not count her data. This disappointed her. This participant countered the issue by testing various locations until the phone counted her activities. These responses suggest that any application that relies on continuous sensing of physical activity must address the issue of phone wearability. One way to significantly improve user experience could be to provide users with a self-report option to correct the number of minutes measured. This feature would be a complement to objective activity monitoring.

Participants who already owned phones carried the smartphones provided in their purse or fanny pack. The main motivation for participants to keep carrying an extra device was their concern that they would appear less active than they actually were. Participants also designed custom harnesses and placed their phones in non-traditional locations such as the knee, pant legs or in socks so that they may count

certain activities that the application was not designed to capture. Participants were disappointed if the phone did not give them credit for movements. Despite the fact that such improvisations were needed, participants didn't consider it inconvenient to do so. This suggests that their motivation to count physical activity superseded the inconvenience of carrying a smartphone in a non-traditional location. Participants did not perceive any changes in their regular phone carrying behavior despite carrying an extra device. All participants felt that it would be more convenient if they could use their existing phones instead of an extra phone.

4) *Social Support and Sharing*: Surveyed participants were ambivalent toward sharing their activity data with peers. One reason for this could be that the application did not make it easy to share or observe data within peers. One participant felt that if there were a real-time component, she would check comparative statistics more often to see how well she was doing. The other participants agreed with this assessment. Participants were ambivalent towards sharing the data from the application with their family but were more favorable toward sharing data with peers. One participant felt that if she was not doing well, she might not share her information. On the other hand, participants also felt that team support could help motivate the people who were not active by reinforcing the idea that every little contribution will help their team. The participant who accumulated the second largest number of minutes felt that by comparing her minutes to the participant with the largest number of minutes, she would have realized that she only needed an extra 15 minutes of activity per day. This would have motivated her to be more active.

An observation was made with three participants during the mid-study check-in sessions: these participants chose the same time to sync their data and thus were in the same room. Unprompted by the coordinator, the participants started talking about their activity summaries their relative progress. During these conversations, they also provided peer support (e.g. "You're doing fine for someone in their 70s") and using the data as a topic to discuss their daily lives. These discussions yielded suggestions and tips that participants used to increase their activity levels (e.g. "I walked the dog for an extra ten minutes to boost my counts"). This suggests that the utility of peer support in a group setting where interested participants can discuss their progress and provide encouragement.

Together, these observations suggest that social features can be a potentially useful tool to increase long-term physical activity levels and also the fun-quotient of such interventions. Potential themes for social support may center around using social circles to provide peer suggestions and support, using team loyalty to encourage activity, peer encouragement and allowing peer comparisons to identify areas of improvement. We aim to explore these dynamics in future extensions of the study.

IV. CONCLUSION

This study provided a proof-of-concept validation of the use of smartphone-based accelerometers to encourage physical activity among older adults, in particular, those between 50 and 80 who have little or no smartphone experience. Participants did not face any major challenges in using the application to track their activities. The main motivations for participants in this study to stay active were loss of

personal health due ageing, the need to stay young and the need to manage chronic conditions. This has implications in tailoring interventions and providing motivational messages to encourage them to stay active.

The value that 24x7 smartphone-based sensors can provide to this population is an easy-to-use and always available companion to check in on their daily lifestyles that allows them to maintain awareness about their physical activity without having to remember what they did. Participants had internal goals that naturally arose from tracking and felt it was important to set personal goals and then have the means to track those goals. If this platform could be coupled with a coaching element from an expert to help participants interpret their data over the long term, the value of the system would be even higher. Participants were willing to bear the inconvenience of physical activity tracking due a phone if it meant that they would obtain a better report of their daily physical activities. However, any application that tracks physical activity using smartphones needs to address the wearability of phones either from within the application itself or through external harnesses to place the phone. Older adults showed lower physical activity.

The main limitations of this study are that it focused on a small population of adults (all women) over a short period of time (three weeks). While it showed a satisfactory initial validation of such a system, it did not supply enough data to make long-term conclusions about the efficacy of smartphones in promoting physical activity. Additionally, the study chose adults who were already motivated to stay active. We would like to expand the scope of this work to a longer duration (16 weeks) study involving 50 participants (both male and female) at various levels of motivation and compare efficacy. The other limitation in this study was that due to the lack of social support features, the potential of peer support to encourage individuals to stay more active was not fully explored. We also plan on providing a self-report feature for people to correct for active minutes that cannot be recorded or can be recorded with great difficulty by the phone. We also plan on providing more options to carry the phone (including lanyards, belt clips or arm bands). Other enhancements include a web dashboard to make data collection and viewing easier.

V. ACKNOWLEDGMENTS

The authors wish to thank Davis Park, Kari Olson, Helen Choo and Julie Santos (Front Porch Center for Technology Innovation and Well being) and Rosa Aguirre, Onyika Lucero and Kim Rathman (OASIS active adults community) for their support and assistance.

REFERENCES

- [1] T. Sugiyama, G. Healy, D. Dunstan, J. Salmon, and N. Owen, "Joint associations of multiple leisure-time sedentary behaviours and physical activity with obesity in australian adults," *International Journal of Behavioral Nutrition and Physical Activity*, vol. 5, no. 1, p. 35, 2008.
- [2] G. Healy, K. Wijndaele, D. Dunstan, J. Shaw, J. Salmon, P. Zimmet, and N. Owen, "Objectively measured sedentary time, physical activity, and metabolic risk the australian diabetes, obesity and lifestyle study (ausdiab)," *Diabetes care*, vol. 31, no. 2, pp. 369–371, 2008.
- [3] F. Hu, T. Li, G. Colditz, W. Willett, and J. Manson, "Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women," *Jama*, vol. 289, no. 14, pp. 1785–1791, 2003.
- [4] M. Reeves, G. Healy, N. Owen, J. Shaw, P. Zimmet, and D. Dunstan, "Joint associations of poor diet quality and prolonged television

- viewing time with abnormal glucose metabolism in australian men and women," *Preventive medicine*, vol. 57, no. 5, pp. 471–476, 2013.
- [5] C. Macera, D. Jones, M. Yore, S. Ham, H. Kohl, C. Kimsey Jr, D. Buchner *et al.*, "Prevalence of physical activity, including lifestyle activities among adults-united states, 2000-2001." *Morbidity and Mortality Weekly Report*, vol. 52, no. 32, pp. 764–766, 2003.
- [6] J. Cylus, M. Hartman, B. Washington, K. Andrews, and A. Catlin, "Pronounced gender and age differences are evident in personal health care spending per person," *Health Affairs*, vol. 30, no. 1, pp. 153–160, 2011.
- [7] S. Pagoto, "The current state of lifestyle intervention implementation research: where do we go next?" *Translational behavioral medicine*, vol. 1, no. 3, pp. 401–405, 2011.
- [8] S. Wilcox, M. Dowda, L. Leviton, J. Bartlett-Prescott, T. Bazzarre, K. Campbell-Voytal, R. Carpenter, C. Castro, D. Dowdy, and A. Dunn, "Active for life: final results from the translation of two physical activity programs," *American Journal of Preventive Medicine*, vol. 35, no. 4, pp. 340–351, 2008.
- [9] A. Smith, "Smartphone adoption and usage," *Pew Research Internet Project*, vol. 15, 2011.
- [10] A. King, E. Hekler, L. Grieco, S. Winter, J. Sheats, M. Buman, B. Banerjee, T. Robinson, and J. Cirimele, "Harnessing different motivational frames via mobile phones to promote daily physical activity and reduce sedentary behavior in aging adults," *PLoS one*, vol. 8, no. 4, p. e62613, 2013.
- [11] M. Fiordelli, N. Diviani, and P. Schulz, "Mapping mhealth research: a decade of evolution," *Journal of medical Internet research*, vol. 15, no. 5, 2013.
- [12] S. Consolvo, D. McDonald, and J. Landay, "Theory-driven design strategies for technologies that support behavior change in everyday life," in *Proceedings of the SIGCHI Conference on Human Factors in Computing Systems*. ACM, 2009, pp. 405–414.
- [13] Y. Fukuoka, J. Komatsu, L. Suarez, E. Vittinghoff, W. Haskell, T. Noorishad, and K. Pham, "The mped randomized controlled clinical trial: applying mobile persuasive technologies to increase physical activity in sedentary women protocol," *BMC public health*, vol. 11, no. 1, p. 933, 2011.
- [14] M. Kirwan, M. Duncan, C. Vandelanotte, and W. Mummery, "Using smartphone technology to monitor physical activity in the 10,000 steps program: a matched case-control trial," *Journal of medical Internet research*, vol. 14, no. 2, 2012.
- [15] S. Munson and S. Consolvo, "Exploring goal-setting, rewards, self-monitoring, and sharing to motivate physical activity," in *Pervasive Computing Technologies for Healthcare (PervasiveHealth), 2012 6th International Conference on*. IEEE, 2012, pp. 25–32.
- [16] C. Manohar, S. Mc Crady, Y. Fujiki, I. Pavlidis, and J. Levine, "Laboratory evaluation of the accuracy of a triaxial accelerometer embedded into a cell phone platform for measuring physical activity," in *2010 Experimental Biology meeting*. FASEB, 2010.
- [17] L. Glynn, P. Hayes, M. Casey, F. Glynn, A. Alvarez-Iglesias, J. Newell, G. ÓLaighin, D. Heaney, and A. Murphy, "Smart move-a smartphone-based intervention to promote physical activity in primary care: study protocol for a randomized controlled trial," *Trials*, vol. 14, no. 1, p. 157, 2013.
- [18] H. Vathsangam, B. Emken, E. Schroeder, D. Spruijt-Metz, and G. Sukhatme, "Energy estimation of treadmill walking using on-body accelerometers and gyroscopes," in *Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*. IEEE, 2010, pp. 6497–6501.
- [19] J. Levine, S. S., and M. Jensen, "Energy expenditure of nonexercise activity," *The American journal of clinical nutrition*, vol. 72, no. 6, pp. 1451–1454, 2000.
- [20] W. Haskell, I. Lee, R. Pate, K. Powell, S. Blair, B. Franklin, C. Macera, G. Heath, P. Thompson, and A. Bauman, "Physical activity and public health: updated recommendation for adults from the american college of sports medicine and the american heart association," *Circulation*, vol. 116, no. 9, p. 1081, 2007.
- [21] C. Wong, C. Khong, and H. Thwaites, "Mobile user interface for seniors: An impact of ageing population on mobile design," *Design Principles & Practice: An International Journal*, vol. 4, no. 4, 2010.