

A Representative Modeling Approach to Sampling Dynamic Ocean Structures

Ryan N. Smith^a, Christoffer Heckman^b, Gabe Sibley^b and M. Ani Hsieh^c

^a *Fort Lewis College, Durango, CO USA*

^b *University of Colorado Boulder, Boulder, CO USA*

^c *Drexel University, Philadelphia, PA USA*

Abstract

The primary goal to this research is to develop a principled approach to sampling in the context of a spatiotemporally dynamic feature. We examine the concepts of registration and association of sparse data in non-static environments with the aim of developing generative models and low-order descriptions to improve localization relative to the dynamic feature, which in turn will improve methods for adaptive sampling. The motivation for this work is in the study of coherent structures within the ocean, e.g., ocean fronts and Lagrangian Coherent Structures (LCS), and how intelligent sampling can help provide an understanding of these complex phenomena. We additionally examine the way that dynamic ocean phenomenon not only provide the features of interest to sample, but also provide the information to generate an environmental representation that drives intelligent adaptation for sampling. The proposed work lies at the intersection of robotics, AI, nonlinear dynamics, perception, network sciences, and ocean science.

1. Introduction

The oceanic environment is both highly dynamic and complex. Yet in this complexity, coherent structures do exist and have recently been shown to hold a wealth of information. Ocean fronts hold the key to understanding many physical and biogeochemical processes, while Lagrangian Coherent Structures (LCS) are essential mechanisms for long-distance transport phenomena. Effective observation and quantification of these coherent structures requires simultaneous measurement of diverse water properties, which must be made rapidly to capture the spatial and temporal variability of multiple

simultaneous interactions. This *cannot be done by traditional oceanographic methods* involving infrequent and sparse measurements from ships, buoys and drifters [1]. In addition, there is no single model that provides an informed view or perception of these or any other ocean feature that enables intelligent sampling in a principled manner. Thus, forecasting where a robot should sample in the immediate future is a challenging task; previously addressed by pre-planned missions designed by oceanographic domain experts. Here, we propose to enable persistent autonomy for intelligent sampling with a novel knowledge transfer approach; learning generative sensor models or low-order descriptions of spatiotemporal phenomenon (such as river plumes) in one environment and leveraging these same algorithms in new situations (such as front tracking in the ocean). Our goal is to *learn how to develop and refine representative models of complex, dynamic phenomena to improve localization and inform adaptive sampling for applications in ocean science.*

2. Technical description of your research

We envision an analogue scenario to that of an international traveler in ocean sampling problems with autonomous underwater vehicles (AUVs). Even without knowing the regional language or having been to the city previously, travelers are able to localize and navigate within built environments (cities) simply by looking at an overhead map [2]. For example, we have seen multiple algal blooms previously, and we have physical and biological models to predict their development, evolution and eventual collapse. However, each algal bloom is unique and must be sampled differently, i.e., no standard sampling path can be implemented on an AUV for effective sampling. There are distinct boundaries and unique regions that exist within every bloom (and similar dynamic features) that could be used for localization and navigation. An added complexity arises in this problem in that instead of a static environment like a city, we are trying to localize and sample within a spatiotemporally dynamic environment.

We generally understand a single feature, e.g., ocean front, algal bloom, etc., well and have basic models and/or representations of them [3, 4, 5, 6, 7, 8]. Sparse data from prior sampling can provide snapshots or representations of certain regions or areas within the feature; environmental niche models [9]. These environmental niche models are very unique, have distinct boundaries and extents, and are the analogues of the memorable objects and places used by humans to navigate and localize in new cities, creating a representation

used to underpin intelligent sampling. To date, there has been minimal progress in utilizing idealized physical models, and specific *memorable* data into navigation/localization within spatiotemporally dynamic phenomena. Additionally, these dynamic processes are not only features to look for or extract when sensing the environment, but they can help us with creating a low-order description or representation of the environment as a whole.

This challenge has motivated our investigation into the utility of combining *a priori* measurements, predictions (from models), and expected *scenes* from data for autonomous localization and sampling in complex and dynamically evolving environments. Specifically, we are focused on the following three fundamental research questions.

1. Can we develop and refine generative sensor models as low-order descriptions of spatiotemporally dynamic features that retain relevant dynamic *scenes* that can be used to predict the structure and motion to inform intelligent sampling of a similar or future feature.
2. How do we validate the accuracy of the model representation, and in what ways can we intelligently design and redesign this representation to increase the accuracy of our predictions of different dynamic phenomena?
3. Can we utilize the generative sensor models to aid and improve localization within spatiotemporally dynamic phenomena in regions where visual-inertial odometry (VIO) techniques are not applicable, e.g., ocean fronts and Lagrangian Coherent Structures (LCS) boundaries in the open ocean.

While many lessons learned in the mapping of and navigation within terrestrial environments translate to analogues underwater, they don't *ALL* translate, and part of the work presented here is to elucidate which ones do and which ones don't due to the need for different sensors and extraction of different features.

3. Key challenges

The capability to develop and refine generative sensor models and predictive representations of a dynamically evolving environment and/or feature, and of interpreting complex meaning from a set of spatial and temporal non-uniform data, will create breakthrough science in the form of new insights

into how intelligent agents generalize and perform adaptive sampling and navigate through complex environments. The ability to learn memorable objects and places in one particular environment and recall those in new environments or situations to accurately predict a representation of a complex environment, represents (up until now) a unique capability of humans (and some animals) to effortlessly extrapolate from past experiences. There are four key challenges in providing an intelligent agent (robots/computer systems/mobile devices) with these predictive tools.

Challenge 1 Learn and improve generative sensor models and low-order representations that represent a spatiotemporally dynamic feature, encoding relationships in observed and predicted data with the capability of *filling in the gaps* for unknown or uncertain regions.

Challenge 2 Utilization of the representative models to improve localization information in domains where GPS and/or visual-inertial odometry is not applicable or available and where dynamic features define the environment.

Challenge 3 Develop the principles necessary to synthesize adaptive sampling plans in the context of a spatiotemporally evolving feature/environment.

Challenge 4 A systematic modeling, decision and assessment approach to model representation design and model quality.

4. The Future

While the ocean presents as deterministic on a long-term scale (and is modeled as such), short-term variability must be modeled using probabilistic models that will guide control and planning algorithm design. As AUVs become less expensive with longer mission times, the deployment of fleets of assets will be realizable, enabling a more synoptic view of ocean processes rather than the sparse subset currently available. The concept of using a cooperative multi-vehicle approach permits the analysis of sensor deployment in a stochastic environment using probabilistic measures for coverage while using minimal on-board power [10]. This will enable models for coherent structures and ocean fronts may be fit for specific flows by amassing data on the movements of sensors in the ambient flow, also informing efficient transport within the medium [11].

The strategy of employing many robot systems has always been one of distributing responsibility, risk, and cost among the various members of a team. In recent years, approaches in the coordination of many robot systems have heavily favored decentralized architectures where individual robots rely solely on local sensing and communication. The arguments for favoring the decentralized approach were the potential for infinite scalability, massive parallelism, and lack of a single point-of-failure. The advent and maturing of cloud computing technologies means that robots are no longer limited by processing constraints and availability of data specifically data that is dependent on proximity. This suggests an entirely different paradigm for coordinating many robot systems where the advantages of both centralized and decentralized strategies can be combined for more effective execution of complex tasks.

Through this research effort, we aim to facilitate a better understanding of the relationship between predictive models and decision, navigation and/or exploitation performance. Additionally, we envision innovation in the ability to utilize predictive models to aid localization, and in turn, drive intelligent sampling and model refinement. Specifically, model synthesis will generate new ways to assimilate data driven process and model forecasts will impact the way stochastic process are projected in both the spatial and temporal dimensions. This will all come together in an integrated framework where ocean flow data can be collected, processed, and assimilated into a high-fidelity ocean model. The high-fidelity model is then continuously updated and used to adapt and improve the sampling and navigation strategies of the AUV team.

- [1] D. Olson, G. Hitchcock, A. Mariano, C. Ashjan, G. Peng, R. Nero, and G. Podesta, "Life on the edge: marine life and fronts," *Oceanography*, vol. 7, pp. 52–60, 1994.
- [2] C. Liu, H.-Y. Shum, and W. Freeman, "Face hallucination: Theory and practice," *International Journal of Computer Vision*, vol. 75, pp. 115 – 134, 10 2007.
- [3] E. D'Asaro, C. Lee, L. Rainville, R. Harcourt, and L. Thomas, "Enhanced turbulence and energy dissipation at ocean fronts.," *Science (New York, N.Y.)*, vol. 332, pp. 318–22, 4 2011.
- [4] A. M. Mancho, E. Hernández-García, D. Small, S. Wiggins, and V. Fernández, "Lagrangian transport through an ocean front in the

- northwestern mediterranean sea,” *Journal of Physical Oceanography*, vol. 38, pp. 1222–1237, 6 2008.
- [5] I. M. Belkin and A. L. Gordon, “Southern ocean fronts from the greenwich meridian to tasmania,” 1996.
- [6] J. Hopkins, P. Challenor, and A. G. P. Shaw, “A new statistical modeling approach to ocean front detection from sst satellite images,” *Journal of Atmospheric and Oceanic Technology*, vol. 27, pp. 173–191, 1 2010.
- [7] J. K. Moore, M. R. Abbott, and J. G. Richman, “Location and dynamics of the antarctic polar front from satellite sea surface temperature data,” 1999.
- [8] J. Hopkins, A. Shaw, and P. Challenor, “The southland front, new zealand: Variability and enso correlations,” *Continental Shelf Research*, vol. 30, pp. 1535–1548, 8 2010.
- [9] J. Elith, S. J. Phillips, T. Hastie, M. Dud’ik, Y. E. Chee, and C. J. Yates, “A statistical explanation of maxent for ecologists,” *Diversity and Distributions*, vol. 17, no. 1, pp. 43–57, 2011.
- [10] C. Heckman, I. Schwartz, and A. Hsieh, “Controlling basin breakout for robots operating in uncertain flow environments,” in *14th International Symposium on Experimental Robotics 2014*, 2014.
- [11] S. C. Shadden, F. Lekien, and J. E. Marsden, “Definition and properties of lagrangian coherent structures from finite-time lyapunov exponents in two-dimensional aperiodic flows,” *Physica D: Nonlinear Phenomena*, vol. 212, no. 3 - 4, pp. 271 – 304, 2005.