

A Neurobiological Perspective on Humanoid Robot Design

Simon F. Giszter, MCP Hahnemann University School of Medicine
Karen A. Moxon and Ilya Rybak, Drexel University
John K. Chapin, State University of New York Downstate

BIOLOGICAL DESIGNS AND NEUROBIOLOGICAL controls continue to inspire technological development. Biology provides working examples and conceptual proofs that push the engineering envelope. The human form and its augmentation are major sources of technological innovation. Engineering helps us interpret biological adaptations observed in nature. Many of our major historical advances, such as tools, telescopes, and writing, are derived from technologies improving action, augmenting perception, and providing cognitive aids to the individual. These advances have increased the speed, power, spatial range, appropriateness, and precision of human actions and perceptions.

During the last two decades, biologically inspired robotics developed into a burgeoning field exploring the ideas of artificial life and adaptive behaviors (see the sidebar, “An expanding field”). It won’t be long before robotic lobsters, cockroaches, flies, lamprey, and tuna enter the commercial market. We hope these efforts lead to the ultimate robot—an autonomous humanoid agent—able to mimic aspects of human action, perception, and cognition in remote or hazardous environments such as deep space or radiation spills. These agents will be invaluable because they will integrate into many tasks without altering existing infrastructures or devices as a result of the human form fac-

tors. In this article, we briefly review recent neurobiological data and ideas that might impact the development of autonomous humanoid robots. We describe motor primitives and exciting results with new neuroprosthetic and neurorobotic technologies. These approaches reveal the inner workings of the most flexible and sophisticated motor controllers in existence. They also provide novel and important insights into biological organization, which can be translated into engineering designs. The framework provides a common language for neuroscientists, engineers, and computer scientists to collaborate in the codevelopment of robotic neuroprosthetic and neural network components. Their joint efforts will improve our understanding of the human brain and ourselves as they propel the design of new control architectures for humanoid robots, new technologies, new biomedical devices, and new interfaces with humanoid robots.

NEUROBIOLOGY AND NEUROENGINEERING CAN OFFER INSIGHTS TO HUMANOID ROBOT DESIGNERS. MOTOR PRIMITIVE CONCEPTS AND MAMMALIAN CORTEX MOTOR CODING PROVIDE A USEFUL FRAMEWORK FOR DESIGNING LEARNING SYSTEMS THAT REPLICATE HUMAN MOTOR SKILLS.

Biological designs for motor control

We would like to emulate animal design features in an autonomous robot. Significant physical design features include energy source and density; sensors and density; and the density, robustness, and flexibility of neuronal axons or “wires.” We are inspired by computational design because of its unrivaled flexibility, fault tolerance, and power to manage vast arrays of sensory information and novel tasks. Biological systems’ physical design falls short of current technologies in the communication speeds between computing elements. Neural axons conduct their digital signals or action potentials at speeds less than 120 meters per second. The maximum rate of action potentials on the axon is low, less than 500 Hz, and there are substantial delays, often of more than 0.3 ms, at the

links of neurons or synapses, where much computation occurs. These biological shortcomings lead to long delays in any neuron control loop.

However, biological systems show that an enormously powerful, robust, and adaptive system can be constructed despite neurons' inherent delays. The human brain is unparalleled for flexible motion control, planning, and abstract cognition. The vast numbers of sensors, the numbers of parallel neurons used to process information, individual neuron's complex processing capabilities, and a highly-evolved architecture all compensate for the delays. These assets let us construct a highly distributed, adaptable, and robust system of computational elements for internal model-based prediction, control, and communication.

Layered redundant competences, partial local autonomy. Perhaps the most striking feature of neurobiological design is a set of layered capabilities. Tasks such as the organism's locomotion or reaching might be represented at many levels in a distributed hierarchy. At each level, the task is represented with different degrees of precision and flexibility. For example, since the late 18th or early 19th century, biologists have recognized that the lowest level of the central nervous system (CNS), the spinal cord, can represent and execute purposeful movements after separation from the brain. For example, a spinal frog—one in which the brain has been surgically severed from the spinal cord—still attempts to remove irritants and uses precisely aimed and adapted trajectories to protect its skin from irritants. This led early psychologists such as William James to attribute consciousness to the spinal frog!

Higher vertebrates, such as humans, have the same organization. Successive surgical lesions of a cat's CNS, which progress from cortex removal to spinal cord isolation, gradually alter locomotion control and organization. Initially, the cat has voluntary locomotion, which gradually deteriorates: from obstinate progression with good balance, to unbalanced spinal locomotion with coordinated forelimbs and hind limbs, to isolated hind limbs and bipedal coordination. The progressive reduction of the cat's CNS shows that the basic competence of locomotor patterning and weight support is organized and available at the CNS's lowest levels. These observations partly inspired Rodney Brooks and his colleagues' subsumption program and

An expanding field

Within the last two decades, researchers' concerted efforts have developed biologically inspired robotics. Laboratories, such as those of Christopher Atkeson, Hans Moravec, Jean-Arcady Meyer, Stewart Wilson, Rodney Brooks, Maja Mataric, Stefan Schaal, Mitsuo Kawato, Patti Maes, David McFarland, Frank Grasso, Joseph Ayers, Roger Quinn and Roy Ritzmann, Randall Beer, Marc Raibert, Daniel Koditschek, Holk Cruse, Stephen DeWeerth, Neville Hogan, Jean-Jacques Slotine, Luc Steels, Edward Colgate, Blake Hannaford, Sebastian Thrun, and Owen Holland have driven the design of autonomous systems, swarms of agents, and biologically inspired control designs and actuators. These scientists have explored the concepts of artificial life and adaptive behaviors in ways too numerous to fully explore here. The hope and promise of this approach is that eventually we will construct an autonomous humanoid agent.

other aspects of nouvelle AI approaches.¹

The idea that some execution functions and details can be delegated to the CNS's lower levels is implicit in the layered hierarchy revealed by the neurobiologist's scalpel. Delegated decisions can execute simultaneously on several levels. For example, the brain of a cat whose leg encounters an obstacle while the cat is pursuing a mouse might not address the detailed correction of that leg's motion. The cat's brain might focus on overall balance and planning total body momentum in anticipation of the pounce or jump to follow. Integrating these layers of controls into an adaptive seamless motion control is a central problem of neurobiological motor control.

Expertise and local autonomy exist at each brain organization level. So, some execution data can be hidden from upper levels. As information flows to the brain from sensors in the limbs, it can be acted upon at the local level or passed up to successive higher levels. This results in significant gating or filtering of sensory input and in reducing information flow to the highest levels of the system, based on behavior and need. Not all brain areas receive the same information. Restricted and abstracted sensory data helps focus the computational resources of each brain area on the most significant features and problems.

In the mammalian CNS, several brain areas cooperate in motion control and planning: the cerebral cortex, the cerebellum, and the basal ganglia. Each area ultimately must exert its controls through direct or indirect actions on the low-level spinal cord and brain stem. Kenji Doya has proposed that quite different types of computation and motor learning occur in these areas.² He argues that the cerebellum controls supervised learning of internal models, the basal ganglia focuses on reinforcement learning, and the cerebral cortex provides unsupervised learning. The use of forward and inverse internal models to execute skilled motions and adapt to new environments is widely accepted as one means of coping with the delays inherent in

the biological motor control loops.^{3,4}

Traditionally, researchers have examined brain areas by recording single cells, often in anesthetized animals. Recently, techniques for multielectrode recording in multiple brain areas in nonanesthetized animals have become available. These techniques let us develop richer data and new experiments and technologies. We believe investigating the motor cortex and cerebellum can be most instructive to humanoid robot designers. The motor cortex is involved with developing novel skills. Motor cortex strokes degrade or disable aspects of skilled voluntary motor performance. Understanding the motor cortex's interaction and cooperation with spinal cord mechanisms in elaborating new skills is important. A humanoid's ability to acquire skills will determine its success or failure in the marketplace. Our research indicates skill development might involve building novel behaviors from a set of spinal parts or *primitives*. We'll describe this spinal "toolbox" in the next sections.

Spinal pattern generators. Neurobiologists have refined our understanding of the spinal cord system. We now know that the spinal system can generate in a feed-forward fashion many neural activity patterns required for motion. *Central pattern generators* are collections of spinal cord neural circuits that produce these patterns;⁵ they are operationally defined. The experimenter records neural activity elicited from a spinal cord, severed from the brain, after the body has been paralyzed—for example, with the Amazon arrow poison curare. The spinal cord exhibits activity that would generate certain behavior patterns if the muscles were not paralyzed. Feedback systems play a role in modulating, phase locking, and controlling the pattern's execution and the resulting motion in the unparalyzed animal. By manipulating sensory feedback systems in paralyzed animals, we can examine some of these controls.

The experiments in which we removed or

altered sensory feedback show a dynamic pattern-organizing system in the low-level spinal cord. Pattern generators might free higher centers from many coordination and synthesis details of multijoint, multilimb motion. This likely lets higher centers control locomotion with lower bandwidth connections, freeing them to focus on global motion control and optimization in relation to higher-order tasks such as hunting, stalking, and fleeing.

Recent experiments with unparalyzed spinalized rats and frogs revealed additional modular motor circuits. Different pattern generators in different behaviors and descending systems might recruit these circuits, which we call *primitives*.⁶⁻⁹ Primitives, which offer features that might impact motion construction and organization at spinal and higher levels, have inspired some robotic designs. They also show striking similarities to independently developed algorithms. Primitives might form a core design feature in the organization of biological motor competence and assist in designing isomorphic humanoid systems.

Force-field primitives

In 1991, we used more sophisticated methods to revisit experiments originally conducted by Luigi Galvani and others in the 18th century. Our research suggested that spinal circuits are partly organized into modules that support and control force patterns in the limb.⁶ We based supporting data on microstimulation of the frogs' surgically isolated spinal cord. We introduced a fine electrode into a frog's spinal cord and stimulated it, activating a small collection of neurons in an approximately 100-micrometer sphere around the electrode tip. By activating the neuron groups, and through their circuit projections to other cells, we elicited and recorded limb forces or movements.

We learned that the spinal cord organizes force fields in response to the electrode stimulation. We examined the forces generated in the limb by holding the limb in various postures while stimulating the same spinal cord site. The forces varied with the limb's position. In this context, a force field maps forces generated in the limb to the limb's configuration and motion. The force-field patterns were convergent, structurally stable through time, and repeatable. By stabilizing a limb posture and actuating the limb to

achieve that posture, the fields provided a specific posture and motion control in the spinal cord. Further exploration showed that the spinal cord produces only a small set of force patterns or *force-field primitives*.⁷⁻¹⁰

Force-field primitives are active force fields that exhibit invariant force vector directions and magnitude balances over time; that is, primitives conserve structures over time.^{7,9} These primitives result from ensemble feed-forward and feedback circuits, which together act as a unit. These circuits assemble the limbs' degrees of freedom (the many muscles and joints), into a biomechanical multijoint force-field unit. Thus, they simplify multijoint control and provide preformed building blocks.

In its most general form, a force-field primitive is a vector function of both position and velocity, modulated by a time varying scalar expressed as

$$F(r, \dot{r}, t) = a(t)\phi_i(r, \dot{r}) \quad (1)$$

$F(r, \dot{r}, t)$ is the observed field derived from a primitive that can be expressed as a scaling $a(t)$ of a fixed field structure $\phi(r, \dot{r})$, a function only of position and velocity.

Our experiments used isometric testing, in which we clamp the limb to prevent movement. So, we can write these descriptions without velocity terms. Under these conditions, at a single limb position, the force vectors generated by a primitive will increase and decrease in magnitude with the primitive's activation-deactivation dynamics. At the same time, force magnitude ratios among the sampled limb positions remain constant; that is, we can express a primitive's time evolution as $a(t) \cdot \phi(r)$, which is the description appearing in most neurobiological papers on primitives.

We only found a few types of force field primitives in the frog's spinal cord. So, we tested how they might be combined using multiple electrodes to explore the combination rules in this limited set. In 80% of our microstimulation trials, we could describe the effect of coactivating primitives simply as the linear sum of the individually activated force fields.¹⁰ In the remaining 20%, we observed a winner-take-all effect in which only one force field of the pair of two co-activated fields was expressed.

For a brief look at related research, see the sidebar, "Relation to other approaches."

Building behaviors from force-field primitives: Theoretical frameworks. In 1988, J.

Edward Colgate and Neville Hogan demonstrated that emulating conservative or passive systems offered specific stability guarantees in limb interactions.¹¹ In 1998, Winfried Lohmiller and Jean-Jacques Slotine showed the stability of classes of nonlinear systems created from linear combinations of stable nonlinear dynamical systems.¹² The frog's spinal primitives have these stability properties, which also are highly desirable in an autonomously acting system and its controller.

From an engineering perspective, the biological systems' fields have interesting properties. Theoretical studies show that summation and magnitude scaling of a few primitives can generate a large range of force-field types. For example, Ferdinando Mussa-Ivaldi has explored ways to synthesize arbitrary smooth control fields from Gaussian primitives.^{4,13} Thus, a force-field collection, like the frog spinal cord collection, could synthesize flexible, adaptable motion control of a multijoint limb. Similar mechanisms have been implemented in robots.¹³⁻¹⁷

The combination of vector superposition in the biological system and these theoretical underpinnings suggests we can design simple principled motion control around dynamic force-field primitives, which could be recruited and combined by sets of planning, patterning, and control systems. Furthermore, this design might offer specific stability guarantees during interactions with different environments.

Primitives in actual spinal behaviors. Mechanisms for combining fields into a motion control for reflexive low-level behaviors could be quite simple. Recent data indicate that the kind of complex goal-directed behaviors organized in the spinal cord (which led William James to posit a spinal soul) correspond to a serial and parallel combinations of primitives.

For example, we examined the precisely targeted multiphase wiping reflex.^{9,18} As we mentioned before, a headless frog removes irritants and avoids obstacles with consistent, accurate trajectories. We showed that the aimed trajectory's formation as well as its rapid online corrections, which circumvent obstacles, can be described very simply as the summation of force-field primitives.

We discovered that, during aimed wiping, spinal frogs utilize a trajectory correction response (TCR) if the wiping limb encounters obstacles in its path. The frog moves its

limb rapidly around the obstacle and reaches its target within a few tens of milliseconds of its normal arrival time. This graceful correction is organized entirely in the spinal cord.

We exploited this TCR when we examined reflex movement construction from force-field primitives and the vector summation hypothesis. We elicited the behavior with an electrical stimulus applied through small clips on the dorsal foot of the target (non-wiping) limb. During unimpeded wiping, the wiping limb moved smoothly until the foot base contacted the opposing target limb. We found that placing an obstacle in the ankle's path elicited the TCR. The frog's response resembled the stepping corrective responses observed in cats, except that we elicited the wiping TCRs with skin contact on the opposite surface of the calf or foot.

The TCR integrated smoothly into the wiping limb's kinematics. The target limb's motion wasn't affected. After the correction, the wiping limb achieved target-limb contact at a posture and time that were remarkably close to the posture and time achieved with no collision. The TCR consisted of a brief hip flexion, which was not observed during the unobstructed movement, and an enhanced knee flexion, which let the ankle clear the obstacle and continue its motion to the target.

How does obstacle detection occur? Our force-field and electromyography data are consistent with the TCR being composed of a superposition of multijoint, force-field motor primitives¹⁸ (see Figure 1). We also observed that the temporal dynamics of muscle group activation and primitives were conserved across the motor program. This property might simplify a control design. Several researchers have proposed that human movement might incorporate a conserved dynamical element.¹⁹ Our data placed this element in the spinal cord at the force-field primitive level. Our frog data supports the idea that all primitives used in a behavior have similar temporal dynamics; that is, they are driven by a common function $a(t)$.¹⁸ We believe that in motion control for the spinal frog, you can express any primitive as

$$\theta(r, \dot{r}, t) = A \cdot a(t) \phi_i(r, \dot{r}) \quad (2)$$

where $a(t)$ resembles a fixed half-cycle oscillation or a fixed impulse response, and A is a scaling factor.

This constrains the method of constructing a time-varying force field $F(r, \dot{r}, t)$ for generating a behavior to the selection of scal-




Figure 1. Panels (a) through (d) show normal frog wipes and correction responses. Panels (a) and (b) show wipes with sensory feedback intact; (c) and (d) show wipes with the ankle skin nerves cut (no skin sensation). In the correction, the frog detects the collision through the skin. In (e), the driving correctional forces are due to an added primitive, which is revealed by a field subtraction of the forces responsible for motion in (d) from the forces responsible for motion in (b). In (f), forces correspond to (b) if the limb was free to move. In (g), forces correspond to (d) if the limb was free to move. Panel (h) shows the extracted corrective force-field primitive obtained as shown in (e). (This figure was redrawn from portions of figures published by William J. Kargo and Simon Giszter.¹⁶)

ings A_i , and the phasing τ_i of the component primitives. In this formulation, we construct the motion control field F as

$$F(r, \dot{r}, t) = \sum_i A_i a(t + \tau_i) \phi_i(r, \dot{r}) \quad (3)$$

Primitives and modularity of biological controls. Our studies and other research support a modular control framework for biological behaviors based on force-field primitives. Moreover, the data suggests that the force-field primitive provides a particular kind of building block for movement. Primitives appear to be separate from the timing systems represented by central pattern generators because they can be modulated or combined into an ongoing pattern without altering the timing systems' other output.

Neurobiologists have speculated about this aspect of neurobiological design. Because amplitude and phase are not separable in nonlinear oscillators, biologists often dismiss a decomposition into separate dynamical timing systems that then recruit the execution systems. However, this decomposition appears likely to be needed in dynamically stable, legged animals. The stable duration of primitives that we have observed might be particularly significant. This stability suggests a quantization or granularity of time scale for motion and action at the spinal level. If this is correct, it has broad implications for how a biological system builds internal models, for motor execution and motor learning, for recovery from injury; for neural prosthetics, and for other neural interfaces. Sim-

Relation to other approaches

Several applications have used force-field- or vector-field-based schemes for navigation or obstacle avoidance,¹ virtual humanoids,² and robotic humanoids.³ Stefan Schaal and Dagmar Sternad used oscillating primitives in seven-degrees-of-freedom arms (see www-slab.usc.edu/research). In our work, when each primitive is recruited, we see something resembling a half cycle, a wavelet-like impulse response, or a time dilated "action potential." The primitive combination scheme emerging from our work is similar to Arthur Prochazka's reflex-chaining and fuzzy-logic schemes of motor pattern construction,⁴ the various pulse step models proposed in biological motor control, and Ferdinando Mussa-Ivaldi's work.⁵

References

1. O. Khatib, "Real Time Obstacle Avoidance of Manipulators and Mobile Robots," *Proc. IEEE Conf. Robotics and Automation*,

IEEE Computer Soc. Press, Los Alamitos, Calif., 1985, pp. 500–505.

2. M.J. Mataric et al., "Behavior-Based Primitives for Articulated Control," *Proc. Fifth Int'l Conf. Soc. for Adaptive Behavior (SAB '98)*, MIT Press, Cambridge, Mass., 1998, pp. 165–170.
3. M.M. Williamson, "Postural Primitives: Interactive Behavior for a Humanoid Robot Arm," *Proc. Fourth Int'l Conf. Soc. for Adaptive Behavior (SAB '96)*, MIT Press, Cambridge, Mass., 1996, pp. 124–135.
4. A. Prochazka, "Proprioceptive Feedback and Movement Regulation," *Handbook of Physiology*, L. Rowell and J. Shepherd, eds., Oxford Univ. Press, New York, 1996, pp. 293–330.
5. F.A. Mussa-Ivaldi, "Modular Features of Motor Control and Learning," *Current Opinion in Neurobiology*, Vol. 9, No. 6, Dec. 1999, pp. 713–717.

ilar internal models based on scaled and phased primitives might be suited to a humanoid robot controller.

The spinal cord circuitry that supports this organization will be particularly interesting for basic science, medical, and biomimetic applications. However, these aspects remain issues for further investigation using both experimental and theoretical methods.

Future directions: Cortical neuroprostheses and motor primitives

We have little information on how spinal primitives are integrated into the processes that underlie mammalian motor skill development. This biological aspect interests biomimetic robot designers because it is critical to building and operating humanoid robots. A 400-pound humanoid robot, designed to load cargo onto ships or aircraft, that behaves like a two-year-old child during its initial bootstrap or tune up might not be commercially successful!

The CNS's layered competences often represent an evolutionary layering (phylogeny). However, their persistence in existing creatures might also represent a developmental sequence (ontogeny) required to progressively construct a distributed adaptive system. A series of critical developmental periods occurs, with critical periods in low-level mechanisms developing before those in upper brain areas. For example, a rat's spinal mechanisms develop and harden into stable structures that support crawling two to five days after birth, which is much earlier than cerebellar or cortical system development two weeks after birth. The low-level systems must be preorganized to sup-

port higher-level organizations.

Developmental processes have inspired several AI research directions, including the organization of cortex spinal cord interactions. After basic connections between the spinal cord and cortex are established (at two weeks in rats), organization and skill in the mammalian cortex and spinal cord can be codeveloped and refined. Rats move rapidly from a waddling gait to precise dynamically stable locomotion. We speculate that motor primitives such as the frog's might act as a bootstrap system, providing a basic competence installed by evolution. We might be able to refine motor skills, "seeding" the vast search space and constraining initial sensorimotor explorations. The African savannah's wildebeest or zebra foal must move with the herd almost immediately. Therefore, basic elements from which efficient locomotory skills can be rapidly elaborated must develop before birth. A commercial humanoid robot will need similar "out of the box" capabilities.

Spinal systems are modified through experiential development and descending control influences. Also, evolutionary and developmental processes have tuned, and possibly optimized, the spinal organization to match mechanical plant structure. Researchers believe that when computational structures that control precise voluntary movements are elaborated in the brain, the spinal cord might also change. Subsequently, the brain likely acts as the spinal cord's adaptive critic, resulting in cooperative changes in task structures. We know that some changes are embedded in the spinal cord. So, cooperative plastic optimization of the energetics and control actions in motion control is probable.

We do not know whether the adult animal preserves the "bootstrap primitives," modi-

fies them, or dissolves them and builds new ones. We know a paraplegic rat or cat can develop strategies for weight-supporting locomotion. However, we don't know whether every paraplegic's spinal cord offers the same competences and control possibilities. If, after development is complete, every spinal cord differs from every other, serious problems arise for determining therapy.

What behavior can we expect if we implement electrical stimulating prostheses⁸ or use a transplant or gene therapy to direct axon regeneration into an adult's spinal cord? This issue is also critical to designing and programming humanoid robots, where repeatable and reliable organization is desirable in system behavior and in the supporting circuitry. A unique humanoid robot that is as difficult to repair after an industrial accident as a human spinal cord would fail commercially.

One of the most significant components of the developmental process for human voluntary movement is how the descending systems learn to represent motion and recruit and control the spinal cord motor apparatus. To understand biological motor primitives for humanoid robots that emulate human manipulation, we must know whether cortical systems (which act directly on the spinal cord) learning new movements in adults use spinal primitives, create new primitives in the spinal cord, or operate completely differently. New techniques using neuroprosthetics are available to address these issues and to incorporate or control humanoid or robotic parts. Ultimately, these techniques might allow us to combine humanoid robot parts with neuroprosthetics to develop specific biomedical applications.

For example, John Chapin's laboratory recently provided the first demonstration that

large-scale multineuron recordings in experimental animals can control robotic devices directly from the brain with sufficient accuracy to obtain a water reward from a dropper.²⁰ This paradigm provides a novel means of examining the motor cortex's organization of voluntary representations and computations. If the cortex is disconnected from its normal motor effectors and given the opportunity to directly control an external robot, how will the brain use this novel environment and interface? Does the cortex seek out regularities in the controlled movement and reinvent its own primitives? Alternatively, does the cortex expect to use certain primitives to start the learning process? Answering these questions might assist humanoid robot design.

NEUROPROSTHETICS AND PRIMITIVE paradigms provide new ways to address the biological layering of controls, hierarchy, and execution. Both methods could translate into useful approaches for designing humanoid controllers. We hope that academic departments and institutions, funding agencies, and industrial liaisons and collaborations will encourage and foster continued interaction of neurobiological and engineering approaches and continued synthesis and cross-fertilization of these disciplines. ■

Acknowledgments

This work was partially funded by National Institutes of Health contract NS34640 to Simon Giszter and NIH contract NS62352 and Office of Naval Research grant N00014-98-1-0679 to John Chapin.

References

1. R.A. Brooks, "From Earwigs to Humans," *Robotics and Autonomous Systems*, Vol. 20, Nos. 2-4, June 1997, pp. 291-304.
2. K. Doya, "What Are the Computations of Cerebellum, Basal Ganglia and Cerebral Cortex?" *Neural Networks*, Vol. 12, 1999, pp. 961-974.
3. M. Kawato, "Internal Models for Motor Control and Trajectory Planning," *Current Opinion in Neurobiology*, Vol. 9, No. 6, Dec. 1999, pp. 718-727.

4. F.A. Mussa-Ivaldi, "Modular Features of Motor Control and Learning," *Current Opinion in Neurobiology*, Vol. 9, No. 6, Dec. 1999, pp. 713-717.
5. O. Kiehn, J. Hounsgaard, and K. Sillar, "Basic Building Blocks of Vertebrate Spinal Central Pattern Generators," *Neurons, Networks, and Motor Behavior*, P. Stein et al., eds., MIT Press, Cambridge, Mass., 1997, pp. 47-60.
6. E. Bizzi, F.A. Mussa-Ivaldi, and S.F. Giszter, "Computations Underlying the Execution of Movement: A Novel Biological Perspective," *Science*, Vol. 253, July 1991, pp. 287-291.
7. S.F. Giszter, F.A. Mussa-Ivaldi, and E. Bizzi, "Convergent Force Field Organized in the Frog's Spinal Cord," *J. Neuroscience*, Vol. 13, No. 2, Feb. 1993, pp. 467-491.
8. S.F. Giszter et al., "Intraspinal Microstimulation: Techniques, Perspectives and Prospects for FES," to be published in *Neural Prostheses for Restoration of Sensory and Motor Function*, J.K. Chapin and K.A. Moxon, eds., CRC Press., Boca Raton, Fla., 2000.
9. S.F. Giszter and W.J. Kargo, "Conserved Temporal Dynamics and Vector Superposition of Primitives in Frog Wiping Reflexes during Spontaneous Extensor Deletions," to be published in *Neurocomputing*, 2000.
10. F.A. Mussa-Ivaldi, S.F. Giszter, and E. Bizzi, "Linear Combinations of Primitives in Vertebrate Motor Control," *Proc. Nat'l Academy of Science*, Vol. 91, Aug. 1994, pp. 7534-7538.
11. J.E. Colgate and N. Hogan, "Robust Control of Dynamically Interacting Systems," *Int'l J. Control*, Vol. 48, 1988, pp. 65-88.
12. W. Lohmiller and J.-J. Slotine, "On Contraction Analysis for Nonlinear Systems," *Automatica*, Vol. 34, 1998, pp. 683-696.
13. F.A. Mussa-Ivaldi, "From Basis Functions to Basis Fields: Vector Field Approximation from Sparse Data," *Biological Cybernetics*, Vol. 67, No. 6, 1992, pp. 479-89.
14. O. Khatib, "Real Time Obstacle Avoidance of Manipulators and Mobile Robots," *Proc. IEEE Conf. Robotics and Automation*, IEEE Computer Soc. Press, Los Alamitos, Calif., 1985, pp. 500-505.
15. M. Cannon and J.-J. Slotine, "Space Frequency Localized Basis Networks for Nonlinear Estimation and Control," *Neurocomputing*, Vol. 9, No. 3, 1995.
16. M.J. Mataric et al., "Behavior-Based Primitives for Articulated Control," *Proc. Fifth Int'l Conf. Soc. for Adaptive Behavior (SAB '98)*, MIT Press, Cambridge, Mass., 1998, pp. 165-170.
17. M.M. Williamson, "Postural Primitives: Interactive Behavior for a Humanoid Robot Arm,"

Proc. Fourth Int'l Conf. Soc. for Adaptive Behavior (SAB '96), MIT Press, Cambridge, Mass., 1996, pp. 124-135.

18. W.J. Kargo and S.F. Giszter, "Rapid Correction of Aimed Movements by Summation of Force-Field Primitives," *J. Neuroscience*, Vol. 20, No. 1, Jan. 2000, pp. 409-426.
19. J.K. Chapin et al., "Realtime Control of a Robot Arm Using Simultaneously Recorded Neurons in the Motor Cortex," *Nature Neuroscience*, Vol. 2, No. 7, July 1999, pp. 664-670.
20. H.I. Krebs et al., "Quantization of Continuous Arm Movements in Humans with Brain Injury," *Proc. Nat'l Academy of Science*, Vol. 96, pp. 4645-4649.

Simon F. Giszter is an assistant professor of neurobiology at the MCP Hahnemann University School of Medicine in Philadelphia and an adjunct associate professor at Drexel University's School of Biomedical Engineering, Science, and Health Systems. He has a BA in Natural Sciences and Zoology from Cambridge University and a PhD from the University of Oregon's Institute of Neuroscience. Contact him at the Dept. of Neurobiology and Anatomy, MCPHU School of Medicine, 3200 Henry Ave., Philadelphia, PA 19129; simon@swamthing.uahs.edu.

Karen A. Moxon is an assistant professor with joint appointments at Drexel University's School of Biomedical Engineering, Science, and Health Systems, and in the Department of Electrical and Computer Engineering. She is an adjunct assistant professor of neurobiology at the MCP Hahnemann University School of Medicine. She has a BS in chemical engineering from the University of Michigan and an MS and PhD in aerospace engineering from the University of Colorado. Contact her at Drexel Univ., School of Biomedical Eng., Science, and Health Systems, 3141 Chestnut St., Philadelphia, PA 19104; karen.moxon@drexel.edu.

Ilya Rybak is a research professor at Drexel University's School of Biomedical Engineering, Science and Health Systems. He has an MS in electrical engineering from Odessa State Polytechnical University, Odessa, Ukraine, and a PhD in biophysics from Leningrad (St. Petersburg) State University. Contact him at Drexel Univ., School of Biomedical Eng., Science, and Health Systems, 3141 Chestnut St., Philadelphia, PA 19104; rybak@drexel.edu.

John K. Chapin is a professor at the Physiology Department of the Health Science Center at the State University of New York Downstate in Brooklyn, New York. He has a BS in biology from Antioch College, Massachusetts, and a PhD in physiology from the University of Rochester, New York. Contact him at the Physiology Dept., Health Science Center, SUNY Downstate, 450 Clarkson Ave., Brooklyn, NY 11203; john_chapin@netmail.hscbklyn.edu.